




Contents lists available at ScienceDirect

Harmful Algae

journal homepage: www.elsevier.com/locate/hal



Inland transport of aerosolized Florida red tide toxins

Barbara Kirkpatrick^a, Richard Pierce^a, Yung Sung Cheng^b, Michael S. Henry^a, Patricia Blum^a, Shannon Osborn^a, Kate Nierenberg^a, Bradley A. Pederson^a, Lora E. Fleming^{c,*}, Andrew Reich^d, Jerome Naar^e, Gary Kirkpatrick^a, Lorraine C. Backer^f, Daniel Baden^e

^a Mote Marine Laboratory, Sarasota, FL 34236, United States^b Lovelace Respiratory Research Institute, Albuquerque, NM 87108, United States^c NSF NIEHS Oceans and Human Health Center, University of Miami Rosenstiel School of Marine and Atmospheric Sciences, Miami, FL 33149, United States^d Florida Department of Health, Tallahassee, FL 32399, United States^e Center for Marine Science Research, University of North Carolina at Wilmington, Wilmington, NC 28409, United States^f National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, GA 30341, United States

ARTICLE INFO

Article history:

Received 17 July 2009

Received in revised form 11 September 2009

Accepted 14 September 2009

Keywords:

Brevetoxins

Harmful algal blooms (HABs)

Red tides

Karenia brevis

Asthma

Air monitoring

ABSTRACT

Florida red tides, an annual event off the west coast of Florida, are caused by the toxic dinoflagellate, *Karenia brevis*. *K. brevis* produces a suite of potent neurotoxins, brevetoxins, which kill fish, sea birds, and marine mammals, as well as sickening humans who consume contaminated shellfish. These toxins become part of the marine aerosol, and can also be inhaled by humans and other animals. Recent studies have demonstrated a significant increase in symptoms and decrease in lung function in asthmatics after only one hour of beach exposure during an onshore Florida red tide bloom.

This study constructed a transect line placing high volume air samplers to measure brevetoxins at sites beginning at the beach, moving approximately 6.4 km inland. One non-exposure and 2 exposure studies, each of 5 days duration, were conducted. No toxins were measured in the air during the non-exposure period. During the 2 exposure periods, the amount of brevetoxins varied considerably by site and by date. Nevertheless, brevetoxins were measured at least 4.2 km from the beach and/or 1.6 km from the coastal shoreline. Therefore, populations sensitive to brevetoxins (such as asthmatics) need to know that leaving the beach may not discontinue their environmental exposure to brevetoxin aerosols.

© 2009 Published by Elsevier Ltd.

1. Introduction

Florida red tides are an annual event off the West Coast of Florida and throughout the Gulf of Mexico, caused by blooms of the toxic dinoflagellate, *Karenia brevis* (*K. brevis*). *K. brevis* produces a suite of potent neurotoxins, brevetoxins, leading to significant fish, sea bird, and marine mammal mortality. In addition, these toxins are filtered and accumulated by bivalves such as clams and oysters; people who consume contaminated shellfish become ill with neurotoxic shellfish poisoning (NSP) (Baden et al., 1995; Keynes, 1979; Kirkpatrick et al., 2003; Watkins et al., 2008). *K. brevis* is an unarmored dinoflagellate that can be easily lysed, releasing brevetoxins into the water. Therefore, *K. brevis* breve-

toxins can be incorporated into the marine aerosol (Pierce et al., 1990). When these toxins are inhaled, brevetoxins can cause upper and lower airway symptoms, particularly in people with chronic respiratory illnesses such as asthma (Fleming et al., 2005a,b, 2007, in press). For many years, the public health message has been that symptoms of exposure to aerosolized Florida red tide would diminish when people left the beach (Steidinger and Baden, 1984; Baden, 1983). However, anecdotal reporting from the community during onshore blooms have suggested that this may not be true for all people, particularly those with underlying respiratory diseases (Kirkpatrick et al., 2006, 2009). Therefore, both exposure and health effects from brevetoxins in marine aerosols need to be measured not only at the beach, but also inland from the beach.

Red tide aerosols are marine aerosols containing aerosolized brevetoxins produced in the air/water interface and they are transported by onshore wind to coastal areas (Cheng et al., 2005; Pierce et al., 1990). Pierce and Cheng have reported on the methods to collect and quantify the amount of airborne toxins at the beach during an onshore Florida red tide (Cheng et al., 2005; Pierce et al., 2005). In our previous beach studies, brevetoxin levels measured in the air have ranged from 1.32 ng/m³ (Cheng et al., 2005) to as high as 93 ng/m³ (Backer et al., 2003). This study explored whether

Abbreviations: PbTx, brevetoxin; cm, centimeter; *K. brevis*, *Karenia brevis*; km, kilometer; LCMS, liquid chromatography mass spectroscopy; µg/L, microgram per liter; ng/m³, nanogram per meter cubed; NSP, neurotoxic shellfish poisoning.

* Corresponding author at: c/o Department of Epidemiology & Public Health, University of Miami School of Medicine, 1120 NW 14th Avenue, Clinical Research Building, Room 1049 (R 669), Miami, FL 33136, United States. Tel.: +1 305 243 5912; fax: +1 305 421 4833.

E-mail address: lfleming@med.miami.edu (L.E. Fleming).

48 brevetoxin exposure continues beyond the beach, and if so, how far
49 inland the brevetoxins travel.

50 2. Methods

51 This study is part of a larger research program investigating the
52 exposures and health effects in humans and animals from
53 aerosolized brevetoxins (Cheng et al., 2005; Fleming et al.,
54 2005a,b, 2007, in press; Pierce et al., 2005). In these "beach studies,"
55 environmental sampling was performed during a 3-day period prior
56 to the beginning of this "transect study" at the beach (Fig. 1, Site A),
57 and then continued throughout the transect study described below.
58 The environmental sampling was performed to establish the
59 presence or absence of a *Karenia* bloom (via *K. brevis* cell count in
60 the water, and brevetoxin levels in the water and air samples) as
61 described in prior study publications (Cheng et al., 2005; Pierce et al.,
62 2005).

63 2.1. Sampler locations and sample collection

64 An inland transect line for the placement of the environmental
65 air samplers was established beginning at Siesta Key Beach
66 (Florida) (Site A), moving eastward/inland to a point approxi-
67 mately 6.4 km from the beach study site (Site F). Six high volume
68 samplers (TE-5000, Tisch Environmental Inc., Village of Cleaves,
69 OH, USA) were placed in locations of convenience to provide
70 electrical power for the samplers and ease of changing filters daily
71 (Fig. 1). The samplers were fitted with 20 cm × 28 cm glass fiber
72 filters (Whatman EPM 2000, Maidstone, England). Air samples
73 were collected continuously from approximately 8 a.m. to 5 p.m.
74 daily for a period of 5 days.

2.2. Toxin analysis

76 As described in Pierce et al. (2005), brevetoxins from the marine
77 aerosol were recovered from the glass fiber filters by extraction for
78 12 h in acetone using a Soxhlet apparatus. The extract was
79 evaporated and transferred to vials for liquid chromatography
80 mass spectroscopy (LC-MS) analysis. Brevetoxin recovery was
81 verified by the addition of standard amounts of brevetoxin PbTx-2
82 and PbTx-3 to each of the three filters, the filters were run for 4 h,
83 and subsequently processed for LC-MS analysis.

84 Brevetoxin analyses were performed by LC-MS using a
85 ThermoFinnigan AqA HPLC/MS obtained from Thermo Electron
86 Corp., Manchester, United Kingdom. The LC consisted of a
87 SpectraSystems, LC Pump P4000, Autosampler AS3000, and a
88 Degasser SCM1000. Mass spectral detection was obtained using
89 an AqA single quad system scanned from 204-1216 AMU with
90 AqA Max 40, and a scan rate of 1.1 scans/s. All analysis was
91 conducted using electrospray with the probe at 3 kV and 250 °C.
92 The column was a Phenomenex Security Guard C-18 guard
93 column with a Phenomenex Luna C-18 5Fm 250 mm × 2 mm
94 Analytical Column. The solvent gradient was composed of
95 acidified (0.3% acetic acid) ACN/H₂O over 40 min. The instrument
96 was calibrated with a standard brevetoxin mix containing PbTx-2
97 and PbTx-3, obtained from the Center for Marine Science, UNC
98 Wilmington, NC, USA.

3. Results

100 Table 1 lists the environmental conditions and water and air
101 brevetoxin measurements at Site A during the beach studies prior
102 to the 5-day transect studies. Table 2 lists the environmental




Fig. 1. Inland transect aerosol sampling locations.

Table 1

Summary of beach studies: environmental data.

Date	Temperature (°C)	Humidity (%)	Wind speed (kph)	Wind direction	<i>K. brevis</i> in water (CELLS/L) ¹	Water toxin (µg/L)	Air Toxin (ng/m ³)
Non-exposure beach study							
October 2004	16.0–27.7	35–91	0–18	Offshore to onshore	<1000–2000	<0.03	<0.05
Exposure beach studies							
February 2005	8.9–23.9	40–97	0–24	Offshore	1,051,000–4,639,000	13–128	0.06–0.09
March 2005	16.7–21.0	14–98	0–19	Onshore	59,000–200,000	2–12	20–38

Table 2
Summary of 5-day inland transect studies: environmental data.

Date	Temperature (°C)	Humidity (%)	Wind speed (kph)	Wind direction	Air toxin (ng/m ³)
Non-exposure 5-day transect					
October 2004	19.4–28.3	50–91	0–26	Offshore to onshore	<0.05
Exposure 5-day transect					
February 2005	8.3–22.8	46–95	0–42	Offshore to onshore	0.05–7.4
March 2005	10.0–22.2	43–95	0–32	Offshore to onshore	0.05–16

conditions and air concentration of brevetoxins during the transect studies.

3.1. Non-exposure (control) beach studies

Non-exposure beach studies were defined when no *K. brevis* cells were measured above detectable levels (<1000 cells/L) in the beach waters at Site A (see Fig. 1) and were conducted from October 16–18, 2004. Additionally, no brevetoxins were detected in the water or aerosol samples above the limits of detection by LC-MS (<0.05 ng/m³ air, 0.03 µg/L water) as shown in Table 1.

3.2. Exposure beach studies

Exposure beach studies were defined as *K. brevis* red tide events (cell counts > 1000 cells/L), and were conducted from February 4 to 6, 2005 and March 11 to 14, 2005. A summary of the cell counts from water samples and the total brevetoxins detected in water and aerosol samples for the beach studies along with *K. brevis* cell counts is presented in Table 1. In the beach study of February 4–6, 2005, the cell counts were above 1 × 10⁶/L, and the brevetoxin concentrations in the water were also high in the range of 13–128 µg/L. However, the air concentrations of brevetoxin were barely detectable (0.06–0.09 ng/m³) because the wind direction during the 3-day beach study periods was offshore. During the exposure beach study of March 11–14, 2005, both the cell counts (59,000–200,000 cells/L) and the water concentrations of brevetoxins (2–12 µg/L) were at moderate levels, but the air concentrations of brevetoxin at Siesta Beach were moderate high at 20–38 ng/m³.

3.3. Inland transect studies

The analysis of total brevetoxin concentrations in the aerosol samples collected during the October 19–23, 2004 inland transect study indicated no detectable levels (<0.05 ng/m³) at any of the sample locations as shown in Table 2. The 5-day non-exposure data are not shown graphically since the levels of brevetoxin were below the limit of detection.




Fig. 2. Inland transect: February 2005.




Fig. 3. Inland transect: March 2005.

The total brevetoxin concentrations in the aerosol samples collected during the February 7–11, and March 15–19, 2005 inland transect studies were detectable, and are shown in Figs. 2 and 3, respectively. Although the highest levels of total brevetoxins were always measured at the beach site (Site A in Fig. 1), brevetoxins were detected as far inland as 4.2 km from the beach (Site E in Fig. 1). The brevetoxin aerosol concentrations in general decreased with increasing distance from the Siesta Beach study site. As the aerosol was transported inland, it was dispersed in the air and some particles were deposited on the ground, resulting in decreased concentration from the beach (Y. Chung, unpublished data).

4. Discussion

The results of this study demonstrate that brevetoxins were transported up to 6.4 km inland from the coastal water source. These data also illustrate the variability from day to day based on cell count (density and/or patchiness of the bloom), wind speed, and wind direction.

The significance of this transect study is that people may still be exposed environmentally to the aerosolized brevetoxins even after they leave the beach. Indeed, if people remain on a barrier island where no point is greater than 1.6 km from a coast, they will most likely be continuing their exposure in any outdoor setting and from all directions if the inland waters also contain *K. brevis* blooms. This is significant as Fleming et al. (2005a,b, 2007, in press) have demonstrated that asthmatics have significantly increased symptoms and decreased lung function after only a 1-h exposure to aerosolized brevetoxins. Furthermore, aerosolized Florida red tide toxins seem to have more chronic effects, including symptoms and pulmonary function changes in asthmatics for several days following the 1-h beach exposure (and/or environmental exposure inland), and increased admissions to emergency rooms for respiratory diseases (including pneumonia, bronchitis, and asthma) during active Florida red tides (Kirkpatrick et al., 2006, 2009, submitted for publication). Currently, the public health message in communities with onshore *Karenia* blooms has only recommended leaving the beach area to avoid aerosol exposure; this message needs to be re-evaluated based on these new findings

174 to take into account the possibility of inland environmental
175 exposure to brevetoxins, particularly for persons with underlying
176 lung diseases such as asthma.

177 4.1. Limitations

178 These findings are based on 1 non-exposure study and 2
179 exposure studies, therefore these data do not reflect the entire span
180 of environmental conditions that may alter the distance the toxins
181 travel. Environmental monitoring over a much longer time period
182 is needed to fully understand the conditions that optimize inland
183 transport, as well as monitoring even further inland. In addition,
184 studies are needed to explore possible exposure to brevetoxins
185 inside of buildings. Cell count data and water sampling and
186 analysis for brevetoxins at the 2 sites located near the water would
187 enhance the assessment of the bloom intensity and location.

188 5. Conclusions

189 This study has documented the transport of brevetoxins during
190 Florida red tide blooms from the beach inland by placing a series of
191 air samplers on an inland transect. This study is the first objective
192 documentation of the transport of *K. brevis* brevetoxins inland.
193 These findings should be considered by healthcare providers,
194 public health officials and policy makers in communities affected
195 by Florida red tide, particularly when making recommendations
196 concerning the avoidance of brevetoxin exposure and health
197 effects for persons with underlying lung disease.

198 Acknowledgements

199 This research was supported by the Centers for Disease Control
200 and Prevention and the Florida Department of Health (Cooperative
201 Agreement: U50/CCU423360-02) as well as by the P01 ES 10594
202 from the National Institute of Environmental Health Sciences. The
203 authors thank the Tropical Breeze Motel (Siesta Key, FL) and the
204 Sarasota residents for allowing air samplers to be placed on their
205 property, and their support of our studies.[SS]

206 References

207 Baden, D.G., 1983. Marine food-borne dinoflagellate toxins. *International Review*
208 *Cytology* 82, 99-150.
209 Baden, D., Fleming, L.E., Bean, J.A., 1995. In: DeWolf, F.A. (Ed.), *Marine Toxins*.
210 *Handbook of Clinical Neurology: Intoxications of the Nervous System Part H*.
211 *Natural Toxins and *Drugs*. Elsevier Press, Amsterdam, pp. 141-175.

Cheng, Y.S., Zhou, Y., Irvin, C.M., Pierce, R., Naar, J., Backer, L.C., Fleming, L.E., 212
Kirkpatrick, B., Baden, D., 2005. Characterization of [marine aerosol for assess-](#)
213 [ment of human exposure to brevetoxins](#). *Environmental Health Perspectives*
214 113 (5), 638-643.
Backer, L.C., Fleming, L.E., Rowan, A., Cheng, Y.-S., Benson, J., Pierce, R., Zaias, J., Bean,
215 J., Bossart, G., Baden, D.G., 2003. Recreational [exposure to aerosolized breve-](#)
216 [toxins during Florida red tide events](#). *Harmful Algae* 2, 19-28.
Fleming, L.E., Kirkpatrick, B., Backer, L.C., Bean, J.A., Wanner, A., Dalpra, D., Tamer, R.,
217 Zaias, J., Cheng, Y.S., Pierce, R., Naar, J., Abraham, W., Clark, R., Zhou, Y., Henry,
218 M.S., Johnson, D., Van de Bogart, G., Bossart, G.D., Harrington, M., Baden, D.G.,
219 2005a. Initial [evaluation of the effects of aerosolized Florida red tide toxins](#)
220 [\(brevetoxins\) in persons with asthma](#). *Environmental Health Perspectives* 113
221 (5), 650-657.
Fleming, L.E., Bean, J.A., Kirkpatrick, B., [Chung, Y.S.](#), Pierce, R., Naar, J., Nierenberg, K.,
222 Backer, L.C., Wanner, A., Reich, A., Zhou, Y., Watkins, S., Henry, M., Zaias, J.,
223 Abraham, W.M., Benson, J., Cassidy, A., Hollenbeck, J., Kirkpatrick, G., Clarke, T.,
224 Baden, D.G. [Exposure and effect assessment of aerosolized red tide toxins](#)
225 [\(brevetoxins\) and asthma](#). *Environmental Health Perspectives*, in press. Q3
226 Fleming, L.E., Kirkpatrick, B., Backer, L.C., Bean, J.A., Wanner, A., Reich, A., Dalpra, D.,
227 Zaias, J., Cheng, Y.S., Pierce, R., Naar, J., Abraham, W.M., Baden, D.G.,
228 2007. [Aerosolized red-tide toxins \(brevetoxins\) and asthma](#). *Chest* 131 (1),
229 187-194.
Fleming, L.E., Backer, L.C., Baden, D.G., 2005b. Overview of [aerosolized Florida red](#)
230 [tide toxins: exposures and effects](#). *Environmental Health Perspectives* 113 (5),
231 618-620.
Keynes, R.D., 1979. Ion channels in the nerve cell membrane. *Scientific American*
232 240, 125-135.
Kirkpatrick, B., Fleming, L.E., Backer, L.C., Bean, J.A., Tamer, R., Kirkpatrick, G., Kane,
233 T., Wanner, A., Dalpra, D., Reich, A., Baden, D.G., 2006. Environmental exposures
234 to Florida red tides: [effects on emergency room respiratory diagnoses admis-](#)
235 [sions](#). *Harmful Algae* 5, 526-533.
Kirkpatrick, B., Fleming, L.E., Bean, J.A., Nierenberg, K., Backer, L.C., Chung, Y.S.,
236 Pierce, R., Reich, A., Naar, J., Zaias, J., Wanner, A., Hollenbeck, J., Baden, D.G.
237 [Aerosolized red tide toxins \(brevetoxins\) and asthma: continued health effects](#)
238 [after 1 hour beach exposure, submitted for publication](#). Q4
239 Kirkpatrick, B., Fleming, L.E., Squicciarini, D., Backer, L.C., Clark, R.C., Abraham, W.,
240 Benson, J., Cheng, Y.S., Johnson, D., Pierce, R., Zaias, J., Bossart, G.D., Baden, D.G.,
241 2003. Literature [review of Florida red tide: implications for human health](#)
242 [effects](#). *Harmful Algae* 3 (2), 99-115.
Kirkpatrick, B., Bean, J.A., Fleming, L.E., Backer, L.C., Akers, R., Wanner, A., Dalpra, D.,
243 Nierenberg, K., Reich, A., Baden, D.G., 2009. [Aerosolized red tide toxins \(bre-](#)
244 [vetoxins\) and asthma: A 10 day follow up after 1 hour acute beach exposure](#). In:
245 Moestrup, et al. (Eds.), *Proceedings of the 12th International Conference on*
246 *Harmful Algae*, International Society for Harmful Algae and Intergovernmental
247 Oceanographic Commission of UNESCO, Copenhagen, pp. 297-299.
Pierce, R.H., Henry, M.S., Proffitt, L.S., Hasbrouck, P.A., 1990. Red tide
248 toxin (brevetoxin) enrichment in marine aerosol. In: Graneli, E., Sund-
249 strom, S., Elder, L., Anderson, D.M.D.M. (Eds.), *Toxic Marine Phytoplankton*.
250 Elsevier Scientific Publishing, New York, pp. 397-402.
Pierce, R., Henry, M.S., Blum, P.C., Hamel, S.L., Kirkpatrick, B., Cheng, Y.S., Zhou, Y.,
251 Irvin, C.M., Naar, J., Weidner, A., Fleming, L.E., Backer, L.C., Baden, D.G., 2005.
252 Brevetoxin composition in water and marine aerosol along a Florida beach:
253 [assessing potential human exposure to marine biotoxins](#). *Harmful Algae* 4 (6),
254 965-972.
Steidinger, K.A., Baden, D.G., 1984. In: *Dinoflagellates*, Spector, D.L. (Eds.), *Toxic*
255 *marine dinoflagellates*. Academy Press, New York, pp. 201-261.
256 Watkins, S.M., Reich, A., Fleming, L.E., Hammond, R., 2008. Neurotoxic [shellfish](#)
257 [poisoning](#). *Marine Drugs* 6 (3), 431-455. 270